

Realization of an Optimally Distinguishable Multi-photon Quantum Superposition.

Francesco De Martini and Fabio Sciarrino

*Dipartimento di Fisica and Istituto Nazionale di Fisica della
Materia, Università "La Sapienza", Roma 00185, Italy*

Abstract

We report the successful generation of an *entangled* multiparticle quantum superposition of pure photon states. They result from a multiple *universal* cloning of a single photon qubit by a high gain, quantum-injected parametric amplifier. The *information preserving* property of the process suggests for these states the name of "*multi-particle qubits*". They are ideal objects for investigating the emergence of the classical world in quantum systems with increasing complexity, the decoherence processes and may allow the practical implementation of the universal 2-qubit logic gates.

Since the golden years of quantum mechanics the interference of classically distinguishable quantum states, first epitomized by the famous "*Schroedinger-Cat*" apologue [1] has been the object of extensive theoretical studies and recognized as a major conceptual paradigm of Physics [2,3]. In modern times the sciences of quantum information (QI) and quantum computation deal precisely with collective processes involving a multiplicity of interfering states, generally mutually entangled and rapidly de-phased by decoherence [4]. In many respects the implementation of this intriguing classical-quantum condition represents today an unsolved problem in spite of recent successful studies carried out mostly with atoms and ions [5,6]. The present work reports on a nearly decoherence-free all optical scheme based on the quantum-injected optical parametric amplification (QI-OPA) of a single photon in a quantum superposition state of polarization (π), i.e. a π – *encoded qubit* [7,8].

Conceptually, the method consists of transferring the well accessible condition of quantum superposition characterizing a single-photon qubit, $N = 1$, to a *mesoscopic*, i.e. multi-photons amplified state $M \gg 1$, here referred to as a "*multi-particle qubit*" (*M-qubit*). In quantum optics this can be done by injecting in the QI-OPA the single-photon *qubit*, $\alpha|H\rangle + \beta|V\rangle$, here expressed in terms of two mutually orthogonal π –states, e.g. horizontal and vertical linear π 's: $|H\rangle$, $|V\rangle$. In virtue of the general *information preserving* property of the OPA, the generated multi-particle state is found to keep the *same* superposition character and the interfering capabilities of the injected qubit, thus realizing the most relevant and striking property of the *M-qubit* condition [8]. Since the present scheme basically realizes the *deterministic* $1 \rightarrow M$ *universal optimal quantum cloning machine* (UOQCM), i.e. able to copy *optimally* any unknown input qubit into $M \gg 1$ copies with the same fidelity, the output state will be necessarily affected by *squeezed-vacuum* (*SV*) noise. This one partially spoils the *exact*, i.e. "classical" distinguishability of the two interfering terms of the superposition. This is but a further manifestation of the quantum "no-cloning theorem" within the present *optimal* process. In a more technical perspective, since any UOQCM can be designed to redistribute *optimally* the initial information into many output channels,

the present scheme is expected to find useful QI applications, e.g. in quantum cryptography [11], [12] and in error correction schemes [13]. In addition, a similar method was successfully adopted recently to realize the first $1 \rightarrow 2$ UOQCM and the first universal quantum NOT-gate [14–16].

Let's refer to the apparatus: Fig. 1. The OPA active element was a nonlinear (NL) crystal slab (BBO: β -barium borate), 1.5 mm thick cut for Type II phase-matching, able to generate by spontaneous parametric down conversion (SPDC) $\vec{\pi}$ – *entangled* pairs of photons. Precisely, the OPA *intrinsic phase* was set as to generate by SPDC *singlet* entangled states on the output modes, a condition assuring the *universality* of the cloning transformation by the QI-OPA scheme [16,7,14]. The photons of each pair were emitted with equal wavelengths (wls) $\lambda = 795nm$ over two spatial modes $-\mathbf{k}_1$ and $-\mathbf{k}_2$ owing to a SPDC process excited by a coherent pump UV field provided by a Ti:Sa Coherent MIRA mode-locked pulsed laser coupled to a Second Harmonic Generator (SHG) and associated with a mode with wavevector (wv) $-\mathbf{k}_p$ and wl $\lambda_p = 397.5nm$. The average UV power was $0.25W$, the *pulse repetition rate* (*rep-r*), $r_{UV} = 7.6 * 10^7 s^{-1}$, and the *coherence time* of each UV pulse as well of the generated single photon pulses were $\tau_{coh} = 140$ fs. The SPDC process implied a 3-wave NL parametric interaction taking place towards the right hand side (r.h.s.) of Fig. 1. The UV pump was back-reflected over the mode $\mathbf{k}_p$ onto the NL crystal by a spherical mirror $\mathbf{M}_p$, with μ -metrically adjustable position $\mathbf{Z}$, thus exciting the main OPA "cloning" process, towards the left hand side (l.h.s.) of Fig.1. By the combined effect of two adjustable optical UV waveplates (wp) ($\lambda/2 + \lambda/4$) acting on the projections of the linear polarization π_p of the UV field on the fixed optical axis of the BBO crystal for the $-\mathbf{k}_p$ and $\mathbf{k}_p$ counter propagating excitation processes, the SPDC excitation was always kept at a very low level while the main OPA amplification could reach *any large intensity*, as we shall see shortly. Precisely, by smartly unbalancing the orientation angles $\vartheta_{\frac{1}{2}}$ and $\vartheta_{\frac{1}{4}}$ of the UV *wp's*, the probability ratio of SPDC emission towards the r.h.s. of Fig.1, of two simultaneous correlated photon pairs and of a single pair was always kept below 3×10^{-2} in any *High Gain* condition. One of the photons of the SPDC emitted

pair, back-reflected by a fixed mirror $\mathbf{M}$, was *re-injected* onto the amplifying NL crystal by the input mode $\mathbf{k}_1$, while the other photon emitted over mode $(-\mathbf{k}_2)$ excited the detector D_T , the *trigger* of the overall *conditional* experiment. The detectors (D) were single-photon SPCM-AQR14. A proper setting of $\mathbf{Z}$ secured the space-time overlapping into the NL crystal of the interacting re-injected pulses with wl's λ_p and λ , and then determined the optimal QI-OPA condition. The time optical walk-off effects due to the crystal birefringence were compensated by inserting in the modes $\mathbf{k}_1$, $\mathbf{k}_2$ and $-\mathbf{k}_2$ three fixed X-cut quartz plates Q and one $\lambda/4$ wp. Before re-injection into the NL crystal, the *pure* input qubit on mode $\mathbf{k}_1$, $|\Psi\rangle_{in} = (\tilde{\alpha}|\Psi\rangle_{in}^{\alpha} + \tilde{\beta}|\Psi\rangle_{in}^{\beta})$, $|\tilde{\alpha}|^2 + |\tilde{\beta}|^2 = 1$, represented by the Bloch sphere shown in Fig. 1, underwent unitary $SU(2)$ *rotation* Φ transformations: $\widehat{U}_i \equiv \exp(-i\sigma_i\Phi/2)$ around the three Cartesian axes $i = x, y, z$ by the combined action of the $\lambda/2$ wp WP_T , of the adjustable *Babinet Compensator* B and of the polarizing beam-splitter PBS_T acting on mode $(-\mathbf{k}_2)$ in virtue of the nonlocality correlating the modes $-\mathbf{k}_1$ and $-\mathbf{k}_2$. These $SU(2)$ transformations are represented in Fig.1 (inset) by circles drawn on the surface of a Bloch-sphere. The main OPA process, i.e. acting towards the l.h.s. of Fig.1 on the injected qubit $|\Psi\rangle_{in}$, was characterized by two different excitation regimes, establishing two corresponding *sizes* of the output *M-qubit*:

A) *Low Gain (LG)* regime, characterized by a low excitation UV energy ($3.5nJ$) per pulse, leading to a small value of the NL parametric "*gain*" : $g = 0.07$.

B) *High Gain (HG)* regime, characterized by a larger value (by a factor ≈ 16) of the *gain*: $g = 1.13$. This condition was attained by a further amplification of the UV "pump" beam by a Ti-Sa regenerative amplifier Coherent-REGA operating at a pulse *rep-r*: $r_{UV} = 2.5 * 10^5 s^{-1}$, with pulse duration: $180fs$: Fig.1.

Let us re-write the state $|\Psi\rangle_{in}$ by expressing the interfering states as Fock product states: $|\Psi\rangle_{in}^{\alpha} = |1\rangle_{1h} |0\rangle_{1v} |0\rangle_{2h} |0\rangle_{2v} \equiv |1, 0, 0, 0\rangle$; $|\Psi\rangle_{in}^{\beta} = |0, 1, 0, 0\rangle$, accounting respectively for 1 photon with horizontal (h) polarization on the input $\mathbf{k}_1$, vacuum state on the input $\mathbf{k}_2$, and 1 photon with vertical (v) π on $\mathbf{k}_1$, vacuum state on the mode $\mathbf{k}_2$. The solution of the QI-OPA dynamical equations is found to be expressed by the *M-qubit*: $|\Psi\rangle \equiv (\tilde{\alpha}|\Psi\rangle^{\alpha} + \tilde{\beta}|\Psi\rangle^{\beta})$,

with

$$|\Psi\rangle^\alpha \equiv \gamma \sum_{i,j=0}^{\infty} (-\Gamma)^i \Gamma^j \sqrt{i+1} |i+1, j, j, i\rangle; \quad |\Psi\rangle^\beta \equiv \gamma \sum_{i,j=0}^{\infty} (-\Gamma)^i \Gamma^j \sqrt{j+1} |i, j+1, j, i\rangle \quad (1)$$

where: $\gamma \equiv C^{-3}$, $C \equiv \cosh g$ and $\Gamma \equiv \tanh g$ [14]. These interfering *entangled*, multi-particle states are *ortho-normal*, i.e. $|\langle \Psi | |\Psi \rangle^j|^2 = \delta_{ij}$ $\{i, j = \alpha, \beta\}$ and *pure*, i.e. fully represented by the operators: $\rho^\alpha = (|\Psi\rangle \langle \Psi|)^\alpha$, $\rho^\beta = (|\Psi\rangle \langle \Psi|)^\beta$. Hence the *pure* state $|\Psi\rangle$ is an entangled quantum superposition of two multi-photon *pure states* and bears the *same* superposition properties of the injected single-photon qubit. The genuine quantum signature of this superposition, namely the *non-definite positive* character of the Wigner function $W(\alpha, \beta)$ of the output $|\Psi\rangle$ in the 8-dimensional complex phase-space $\alpha_j, \beta_j, j = 1, 2$, was confirmed by a previous theoretical analysis [7]. For the sake of completeness, consider the overall output density operator $\rho \equiv |\Psi\rangle \langle \Psi|$ and his *mixed-state* reductions over the $\vec{\pi}$ -*vector* spaces relative to the modes $\mathbf{k}_1$ and $\mathbf{k}_2$: $\rho_1 = Tr_2 \rho$; $\rho_2 = Tr_1 \rho$. These ones may be expanded as a weighted superpositions of p -*square* matrices of order $p = (n+2)$, the relative weight Γ^2 of each two successive matrices being determined by the parametric *gain*. Note that Γ^2 approaches asymptotically the unit value for large g . In turn, the p -*square* matrices may be expressed as sum of 2×2 matrices as shown by the following expressions:

$$\rho_1 = \gamma^2 \sum_{n=0}^{\infty} \Gamma^{2n} \times \sum_{i=0}^n \begin{bmatrix} |\tilde{\beta}|^2 (n-i+1) & \tilde{\alpha}^* \tilde{\beta} \sqrt{(i+1)(n-i+1)} \\ \tilde{\alpha} \tilde{\beta}^* \sqrt{(i+1)(n-i+1)} & |\tilde{\alpha}|^2 (i+1) \end{bmatrix} \quad (2)$$

written in terms of the Fock basis: $\{|i\rangle_{1h} |n-i+1\rangle_{1v}; |i+1\rangle_{1h} |n-i\rangle_{1v}\}$. Correspondingly:

$$\rho_2 = \gamma^2 \sum_{n=0}^{\infty} \Gamma^{2n} \times \sum_{i=0}^{n+1} \begin{bmatrix} |\tilde{\beta}|^2 (n-i+1) & -\tilde{\alpha}^* \tilde{\beta} \sqrt{(n-i+1)i} \\ -\tilde{\alpha} \tilde{\beta}^* \sqrt{(n-i+1)i} & |\tilde{\alpha}|^2 i \end{bmatrix} \quad (3)$$

in terms of the Fock basis: $\{|n-i\rangle_{2h} |i\rangle_{2v}; |n-i+1\rangle_{2h} |i-1\rangle_{2v}\}$. Interestingly, the value n appearing in Eqs. 2, 3 coincides with the number of photon pairs generated by the QI-OPA amplification. Note the *non-diagonal* character of these matrices implying the quantum superposition property of the overall state. Furthermore the interfering states possess the *nonlocal* property of all entangled quantum systems. It is worth noting that the

Von Neumann entropies $S(\rho_j) = -\text{Tr}(\rho_j \log_2 \rho_j)$, $j = 1, 2$ are equal, thus implying the same *degree of mixedness* on the 2 output channels: $S(\rho_1) = S(\rho_2)$ [14,17].

The actual $|\Psi\rangle$ experimental detection of the output states could be undertaken by measurements taken either (a) on the injection, i.e. cloning mode $\mathbf{k}_1$, or (b) on the *anticloning* mode $\mathbf{k}_2$, or (c) on both output modes. The option (b) was selected since there the field on mode $\mathbf{k}_2$ is not affected by the input qubit in absence of the QI-OPA action and then any registered *interference* effect is by itself an unambiguous demonstration of the *M-qubit* condition. This condition was carefully verified experimentally. In addition, it was checked that only *non interfering* SV-noise was detected on the output mode $\mathbf{k}_2$, in absence of the injection qubit $|\Psi\rangle_{in}$.

Since the 1st-order interference property of any quantum object is generally expressed by the 1st-order correlation-function $G^{(1)}$ [18], this quantity was measured by two detectors D_2 and D_{2^*} coupled to the mutually orthogonal fields $\hat{c}_j(t)$, $j = 1, 2$ emerging from the polarizing beam-splitter PBS_2 inserted in the output $\mathbf{k}_2$: $G_{2j}^{(1)} \equiv \langle \Psi | \hat{N}_j(t) | \Psi \rangle$. In the present context the $G^{(1)}$ are expressed by ensemble averages of the number operators $\hat{N}_j(t) \equiv \hat{c}_j^\dagger(t) \hat{c}_j(t)$, written in terms of the detected fields: $\hat{c}_1(t) \equiv 2^{-\frac{1}{2}}[\hat{b}_h(t) + \hat{b}_v(t)]$, $\hat{c}_2(t) \equiv 2^{-\frac{1}{2}}[\hat{b}_h(t) - \hat{b}_v(t)]$. There the $\hat{b}_i(t)$, $i = h, v$, defined on the QI-OPA output $\mathbf{k}_2$, underwent a $45^\circ\pi$ -rotation by the $\lambda/2$ -wp WP_2 before injection into PBS_2 : Fig. 1. The same operations transformed the basis $\{h, v\}$ into $\{H, V\}$. By expressing the injected qubit as: $|\Psi\rangle_{in} \equiv (\alpha |\Psi\rangle_{in}^\alpha + \beta e^{i\varphi} |\Psi\rangle_{in}^\beta)$ with α and β real numbers, the $G_{2j}^{(1)}$ could be given as: $G_{2H}^{(1)} = \bar{n} + \frac{1}{2}\bar{n}[1 + 2\alpha\beta \cos \varphi]$; $G_{2V}^{(1)} = \bar{n} + \frac{1}{2}\bar{n}[1 - 2\alpha\beta \cos \varphi]$, showing the superposition character of the output average field per mode with respect to α, β and φ , where $\bar{n} \equiv \sinh^2 g$. By comparison of this result with the corresponding *non interfering* averages $G_{21,vac}^{(1)} = G_{22,vac}^{(1)} = \bar{n}$ taken in absence of the injected qubit, i.e. over the *input vacuum state*, the signal-to-noise ratio for the *M-qubit* detection: was found: $S/N = 2$, for $\varphi = 0$ and $\alpha = \beta = 2^{-\frac{1}{2}}$.

The above result immediately suggests, as a demonstration of the 1st-order π -interference of the *M-qubit*, to draw the *fringe* patterns expressing the quantity: $\Delta G_2^{(1)} \equiv$

$(G_{2H}^{(1)} - G_{2V}^{(1)})$ and bearing the fringe "visibility": $\mathcal{V} = (G_{2\max}^{(1)} - G_{2\min}^{(1)}) / (G_{2\max}^{(1)} + G_{2\min}^{(1)}) = 2\alpha\beta/3$. The experimental values of $G^{(1)}$ were obtained by coincidence measurements involving the couples of detectors: $[D_2 D_T]$ and $[D_2^* D_T]$. Note that the fringe patterns, reported in Fig. 2 and the corresponding values of $\mathcal{V}$ were strongly affected by the superposition character of the injection qubit and by the corresponding $SU(2)$ transformations expressed by the closed paths drawn on the Bloch sphere: Fig.1 (inset). In order to show more clearly the quantum efficiency of the M -qubit for the two "gain" regimes, the *relative coincidence rates*, $\xi = r_{SC}/r_{UV}$ were reported in Fig. 2, where r_{SC} expressed the values of $\Delta G_2^{(1)} \equiv (G_{2H}^{(1)} - G_{2V}^{(1)})$. The experimental best-fit curves drawn in Figure 2 were found to reproduce very closely the quantum theoretical results.

In the LG excitation condition the *average* number of quantum-interfering photon pairs was $\overline{N} \approx 0.009$ while the HG condition it was $\overline{N} \approx 4$, corresponding to the realization of a much "fatter" M -qubit. The value of *average* number of generated pairs was $\overline{N} = 3\overline{n}$ in virtue of the stimulated emission process. They were estimated by accounting for the overall *quantum efficiency* of the D detectors : $QE \approx 18\%$. Calibrated attenuation filters placed in front of the D 's assured the condition of single-photon detection in the HG regime. By the Eqs. 2, 3 a straightforward evaluation of the output photon-pair distribution $p(n)$ could be carried out. By this one, the probability of generating a number of pairs $N \geq 2\overline{N} = 8$, i.e. a quantum superposition in excess of 16 photons, was found to be $P(\overline{N}) \equiv \sum_{k=2\overline{N}}^{\infty} p(k) \approx 14\%$. By moving the injected state with $\alpha = \beta$ over the Bloch sphere, a visibility $V_{th} = 33\%$ of the fringe-pattern was expected from theory. Due to experimental imperfections, mostly due to the mode selection before detection, we have measured $V \approx 4\%$ (4%) with a 2-coincidence detection measurement scheme and $V \approx 25\%$ (7%) with a 4-coincidence scheme respectively for the LG and HG regimes. The results above are expected to be *linearly scaled* by adoption of a more efficient NL crystal and of a more powerful UV source.

Unlike other systems involving atoms or ions [6,5] a limitation of our method consists of the impossibility of controlling the "distance", d on the *phase-space* of the interfering

multi-particle states by means of the QI-OPA parameters. This condition, implied by the results of the Wigner function analysis [7] can be expressed here by the Hilbert-Schmidt (d) of the interfering states: $d(\rho^\alpha; \rho^\beta) = \text{Tr} [(\rho^\alpha - \rho^\beta)^2]$ [19]. This quantity, not affected by the amplification process, is expressed in our case by: $d(\rho^\alpha; \rho^\beta)_{in} = d(\rho^\alpha; \rho^\beta)_{out} = 2$. However, as a lucky counterpart of this condition, the de-coherence of our system can be deemed generally irrelevant as it is mostly determined by the stray reflection losses on the surfaces of the optical components of the apparatus. Accordingly, a number of photons in the range $(10^2 \div 10^3)$ could be made simultaneously excited in quantum superposition without sizable decoherence effects: this is but one of the commendable qualities of the device. Of course, the injection into the OPA of a many particle state will result in a better S/N ratio. The M -qubit condition attained by injection into OPA of a symmetrized 2 photon state is presently in progress in our Laboratory.

In summary, we have realized the interference of classically distinguishable *multi-particle*, *orthonormal*, *pure* quantum states. These characteristics indeed express the peculiar properties of the system originally proposed by Schroedinger [1] with a relevant additional feature: the *entanglement* of the interfering states. The adoption of the OPA to *clone universally* a single qubit in a large gain regime witnesses the scalability of the information preserving property of this device in the quantum domain. Furthermore and very important, unlike other systems involving atoms, e.g. in hardly accessible optical traps or superconducting cavities [6,5], the *M-qubit* generated by our system is directly accessible to measurement or to further exploitation, e.g. by injection into a QI *gate*. In a more conceptual perspective, the present realization could open a new trend of studies on the persistence of the validity of several crucial laws of quantum mechanics for entangled mixed-state systems of increasing complexity [5] and on the violation of Bell-type inequalities in the multi-particles regime [20]. This work has been supported by the FET European Network on QI and Communication (Contract IST-2000-29681: ATESIT), by INFN (PRA "CLON") and by MIUR (COFIN 2002). We thank V. Buzek, S. Popescu, for stimulating discussions and D. Pelliccia for early experimental collaboration.

REFERENCES


- [1] E. Schroedinger, *Naturwissenschaften* **23**, 807-823 (1935).
- [2] A. Caldeira, A. Leggett, *Physica A* **121**, 587 (1983).
- [3] A. Leggett, A. Garg, *Phys. Rev. Lett.* **54**, 857 (1985).
- [4] W. Zurek, *Physics Today*, October 1991, pag.36.
- [5] M. Brune, *et al.*, *Phys. Rev. Lett.* **77**, 4887 (1996), A. Auffeves, *et al.* *Phys. Rev. Lett.* **91**, 230405 (2003), M.W. Noel, *et al.* *Phys. Rev. Lett.* **77**, 1913 (1996).
- [6] C. Monroe, *et al.*, *Science* **272**, 1131 (1996); D. Leibfried, *et al.* *Science* **304**, 1476 (2004), C.F. Roos, *et al.* *Science* **304**, 1478 (2004).
- [7] F. De Martini, *Phys. Rev. Lett.* **81**, 2842 (1998). The most relevant results of the present experiment are reported by F. De Martini, M. Ricci, F. Sciarrino in: "Quantum State Estimation", ed. by M.G.A. Paris and J. Rehacek (Springer Verlag, Berlin 2004).
- [8] F. De Martini, G. Di Giuseppe and S. Padua, *Phys. Rev. Lett.* **87**, 150401 (2001). There the double-injection did not allow a direct detection of the interference.
- [9] W.K. Wootters, and W.H. Zurek, *Nature (London)* **299**, 802 (1982).
- [10] W. Dur, C. Simon, and J.I. Cirac, *Phys. Rev. Lett.* **89**, 210402 (2002).
- [11] H. Bechmann-Pasquinucci and N. Gisin, *Phys. Rev. A* **59**, 4238 (1999).
- [12] E.F. Galvão, and L. Hardy, *Phys. Rev. A* **62**, 022301 (2000).
- [13] D. Bruß, J. Calsamiglia, and N. Lütkenhaus, *Phys. Rev. A* **63**, 042308 (2001).
- [14] F. De Martini, V. Buzek, F. Sciarrino, and C. Sias, *Nature (London)* **419**, 815 (2002).
- [15] F. De Martini, D. Pelliccia and F. Sciarrino, *Phys. Rev. Lett.* **92**, 067901 (2004).
- [16] A. Lamas-Linares, C. Simon, J.C. Howell, and D. Bouwmeester, *Science* **296**, 712 (2002).

- [17] C.H. Bennett, D.P. Di Vincenzo, J.A. Smolin and W.K. Wootters, Phys. Rev. A **54**, 3824 (1996).
- [18] D. Walls, G. Milburn, *Quantum Optics* (Springer, N.Y. 1994), chs.5, 16.
- [19] M. Nielsen and I. Chuang, *Quantum Computation and Information* (Cambridge, 2000) Ch9.
- [20] M.D. Reid, W.L. Munro, and F. De Martini, Phys. Rev. A **66**, 033801 (2002).

Figure Captions


Figure 1. Layout of the *quantum-injected OPA* apparatus. The device **A** represents the Regenerative Ti:Sa laser Amplifier to attain the High Gain (HG) dynamical condition. IN-SET: Bloch sphere representation of the $SU(2)$ unitary transformations applied to the input qubit.

Figure 2. Multi-photon interference fringe patterns expressed by the *relative coincidence rate*, $\xi = \eta(r_{SC}/r_{UV})$ proportional to the ratio of the 2-detector coincidence rates expressing the quantity: $\Delta G_2^{(1)} \equiv (G_{2H}^{(1)} - G_{2V}^{(1)})$ and of the repetition rate of the UV pump excitation in *LG* and *HG* regimes ($\eta = 7.6 \times 10^7$). The patterns correspond to the $SU(2)$ transformations shown in Fig.1 (inset) and affecting the injected qubit. $G^{(1)}$ are the *1st – order correlation functions*.



Relative coincidence rates

Low Gain



High Gain

